



● アトロピン-0.01% : 2年に渡る近視進行度 → 平均 -0.49D

▲ 薬効成分なし(プラセボ) : 2年に渡る近視進行度 → 平均 -1.20D

※ Chia A, Chua WH, Wen L, Fong A, Goon YY, Tan D. Atropine for the treatment of childhood myopia: changes after stopping atropine 0.01%, 0.1% and 0.5%. Am J Ophthalmol. 2014 Feb;157(2):451-457.e1. doi: 10.1016/j.ajo.2013.09.020.
 Chia A, Chua WH, Cheung YB, Wong WL, Lingham A, Fong A, Tan D. Atropine for the treatment of childhood myopia: safety and efficacy of 0.5%, 0.1%, and 0.01% doses (Atropine for the Treatment of Myopia 2). Ophthalmology. 2012 Feb;119(2):347-54. doi: 10.1016/j.ophtha.2011.07.031.
 Tong L, Huang XL, Koh AL, Zhang X, Tan DT, Chua WH. Atropine for the treatment of childhood myopia: effect on myopia progression after cessation of atropine. Ophthalmology. 2009 Mar;116(3):572-9. doi: 10.1016/j.ophtha.2008.10.020. Epub 2009 Jan 22.
 Chua WH, Balakrishnan V, Chan YH, Tong L, Ling Y, Quah BL, Tan D. Atropine for the treatment of childhood myopia. Ophthalmology. 2006 Dec;113(12):2285-91. Epub 2006 Sep 25.